

C1  
cancel

the present invention, in particular embodiments, also relates to transgenic versions of the claimed hybrid maize line 31R88.

In the Claims

Please amend claims 6, 8, 11, 15, 19, 21, 24, 28 and 32 as follows:

6. (Twice Amended)

C2

The tissue culture according to claim 5, the cells or protoplasts of said cells having been isolated from a tissue selected from the group consisting of leaves, pollen, embryos, roots, root tips, anthers, silks, flowers, kernels, ears, cobs, husks, and stalks.

8. (Twice Amended)

Sub E4  
C3

The maize plant of claim 2 wherein said maize plant further comprises a genetic factor conferring male sterility.

11. (Twice Amended)

C4

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 2, wherein said maize plant has derived at least 50% of its alleles from 31R88 and is capable of expressing a combination of at least two 31R88 traits selected from the group consisting of: a relative maturity of 119 based on the Comparative Relative Maturity Rating System for harvest moisture of grain, yield potential under low to moderate yield environments, stalk lodging resistance, root lodging resistance, Staygreen, drought tolerance, resistance to Gray Leaf Spot, resistance to common rust, resistance to Southern Leaf Blight, and brittle stalk resistance.

15. (Twice Amended)

C5

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 21, wherein said maize plant has derived at least 50% of its alleles from 31R88 and is capable of expressing a combination of at least two 31R88 traits selected from the group consisting of: a relative maturity of 119 based on the Comparative Relative Maturity Rating System for harvest moisture of grain, yield potential under low to moderate yield

C5  
consolid

environments, stalk lodging resistance, root lodging resistance, Staygreen, drought tolerance, resistance to Gray Leaf Spot, resistance to common rust, resistance to Southern Leaf Blight, and brittle stalk resistance.

19. (Twice Amended)

C6

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 16, wherein said maize plant has derived at least 50% of its alleles from 31R88 and is capable of expressing a combination of at least two 31R88 traits selected from the group consisting of: a relative maturity of 119 based on the Comparative Relative Maturity Rating System for harvest moisture of grain, yield potential under low to moderate yield environments, stalk lodging resistance, root lodging resistance, Staygreen, drought tolerance, resistance to Gray Leaf Spot, resistance to common rust, resistance to Southern Leaf Blight, and brittle stalk resistance.

Sub E6  
C7

21. (Twice Amended)

The maize plant of claim 2 wherein said maize plant further comprises a genetic factor conferring male sterility.

24. (Twice Amended)

C8

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 24, wherein said maize plant has derived at least 50% of its alleles from 31R88 and is capable of expressing a combination of at least two 31R88 traits selected from the group consisting of: a relative maturity of 119 based on the Comparative Relative Maturity Rating System for harvest moisture of grain, yield potential under low to moderate yield environments, stalk lodging resistance, root lodging resistance, Staygreen, drought tolerance, resistance to Gray Leaf Spot, resistance to common rust, resistance to Southern Leaf Blight, and brittle stalk resistance.

## 28. (Twice Amended)

C9

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 25, wherein said maize plant has derived at least 50% of its alleles from 31R88 and is capable of expressing a combination of at least two 31R88 traits selected from the group consisting of: a relative maturity of 119 based on the Comparative Relative Maturity Rating System for harvest moisture of grain, yield potential under low to moderate yield environments, stalk lodging resistance, root lodging resistance, Staygreen, drought tolerance, resistance to Gray Leaf Spot, resistance to common rust, resistance to Southern Leaf Blight, and brittle stalk resistance.

## 32. (Twice Amended)

C10

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 29, wherein said maize plant has derived at least 50% of its alleles from 31R88 and is capable of expressing a combination of at least two 31R88 traits selected from the group consisting of: a relative maturity of 119 based on the Comparative Relative Maturity Rating System for harvest moisture of grain, yield potential under low to moderate yield environments, stalk lodging resistance, root lodging resistance, Staygreen, drought tolerance, resistance to Gray Leaf Spot, resistance to common rust, resistance to Southern Leaf Blight, and brittle stalk resistance.

Please add new claims 33 - 42 as follows:

## 33. (New)

C11

A method of making a hybrid maize plant designated 31R88 comprising:  
crossing an inbred maize plant GE528776, deposited as PTA-4282 with a second inbred maize plant GE492452, deposited as PTA-4278; and  
developing from the cross a hybrid maize plant representative seed of which having been deposited under ATCC Accession Number PTA-4272.

Sub E 8

34. (New)

A method of making an inbred maize plant comprising:  
obtaining the plant of claim 2 and  
applying double haploid methods to obtain a plant that is homozygous at essentially every locus,  
said plant having received all of its alleles from maize hybrid plant 31R88.

35. (New)

A method for producing an 31R88 progeny maize plant comprising:

- (a) growing the plant of claim 2, and obtaining self or sib pollinated seed therefrom;  
and  
(b) producing successive filial generations to obtain a 31R88 progeny maize plant.

36. (New)

A maize plant produced by the method of claim 35, said maize plant having received all  
of its alleles from hybrid maize plant 31R88.

37. (New)

A method for producing a population of 31R88 progeny maize plants comprising:

- (a) obtaining a first generation progeny maize seed produced by crossing the maize  
plant of claim 2 with a second maize plant;  
(b) growing said first generation progeny maize seed to produce F<sub>1</sub> generation maize  
plants and obtaining self-pollinated seed from said F<sub>1</sub> generation maize plants; and  
(c) repeating the steps of growing and harvesting successive filial generations to  
obtain a population of 31R88 progeny maize plants.

38. (New)

The population of 31R88 progeny maize plants produced by the method of claim 37, said  
population, on average, deriving at least 50% of its alleles from 31R88.

39. (New)

A 31R88 maize plant selected from the population of 31R88 progeny maize plants produced by the method of claim 37, said maize plant deriving at least 50% of its alleles from 31R88.

40. (New)

The method of claim 37, further comprising applying double haploid methods to said F<sub>1</sub> generation maize plant or to a successive filial generation thereof.

41. (New)

A method of producing a male sterile maize plant comprising transforming the maize plant of claim 2 with a genetic factor conferring male sterility.

42. (New)

The method of claim 41 wherein a male sterile maize plant is produced.